

Editorial Office: International Institute of Anticancer Research,
DELINASIOS G.J & CO G.P., Kapandriti, POB 22, Attiki 19014,
Greece. Fax: 0030-22950-53389; Tel: 0030-22950-52945
e-mail: journals@iiar-anticancer.org

Dear Sir/Madam:

Enclosed are the galley proofs of your article for IN VIVO.

We would like to call your attention to the following:

1. Please read thoroughly, correct, and return the proofs to the Editorial Office **within 24 hours**.
2. Proofs should be returned preferably **by e-mail** or **fax**. Delays in the return of these proofs will necessitate the publication of your paper in a later issue of the journal.
3. Please read the entire manuscript carefully to verify that no changes in meaning have been introduced into the text through language improvements or editorial corrections.
4. Corrections should be limited to typographical errors.
5. Should you require reprints, PDF file, online open access, issues or special author rate subscriptions, please fill the attached reprint order form.
6. If you opt for online open access publication of your paper, it will instantly upon publication be available to read and reproduce free of charge.
7. Should you require information about your article (publication date, volume, page numbers, etc) please call: +30-22950-52945 or send an e-mail to journals@iiar-anticancer.org.
8. Please provide your complete address (not P.O.B.), telephone and fax numbers for the delivery of reprints and issues.
9. Please feel free to contact us with any queries that you may have (Tel./Fax: +30-22950-53389 or +30-22950-52945, e-mail: journals@iiar-anticancer.org).

Thank you for taking the time to study these guidelines.

I greatly appreciate your cooperation and your valuable contribution to this journal.

Yours sincerely,



J.G. Delinasios
Managing Editor

Enclosures



Assessment of Oxidative Stress in Septic and Obese Patients Using Markers of Oxidation-reduction Potential

YPATIOS SPANIDIS¹, NIKOLAOS GOUTZOURELAS¹, DIMITRIOS STAGOS¹, ANASTASIA S. KOLYVA², CHARALAMBOS A. GOGOS², DAVID BAR-OR^{3,4,5,6} and DIMITRIOS KOURETAS¹

¹Department of Biochemistry and Biotechnology, University of Thessaly, Larissa, Greece;

²Division of Infectious Diseases, Department of Internal Medicine, Patras University Hospital, Rion-Patras, Greece;

³Trauma Research Department, St. Anthony Hospital, Lakewood, CO, U.S.A.;

⁴Trauma Research Department, Swedish Medical Center, Englewood, CO, U.S.A.;

⁵Trauma Research Department, Medical Center of Plano, Plano, TX, U.S.A.;

⁶Luoxis Diagnostics, Inc., Englewood, CO, U.S.A.

Abstract. *Background/Aim: The novel static (sORP) and capacity (cORP) oxidation-reduction potential markers were examined for assessing oxidative stress in plasma of patients with sepsis. Moreover, the possible effect of obesity-induced oxidative stress on patients with sepsis was investigated. Materials and Methods: sORP and cORP markers, as well as the conventional oxidative stress biomarkers total antioxidant capacity (TAC), thiobarbituric acid-reactive substances (TBARS) and protein carbonyls (CARB), were assessed in plasma. Results: sORP marker was increased significantly in the sepsis group, while cORP was significantly lower compared to the control group, indicating oxidative stress. Furthermore, in patients with sepsis, TAC was significantly lower compared to control group. However, obesity had no effect on sORP, cORP and TAC in patients with sepsis, although it increased levels of CARB and TBARS. Conclusion: The present results suggest for the first time that ORP markers could be used for assessing oxidative stress in patients with sepsis.*

Reactive oxygen species (ROS) are products of normal metabolism (1). ROS include free radicals such as superoxide radical ($O_2^{\bullet-}$), hydroxyl radical (OH^{\bullet}), peroxy radical (RO_2^{\bullet}), as well as nonradical species such as hydrogen peroxide (1). Some ROS are necessary for

physiological processes such as phagocytosis (2), intracellular signaling (1), cell proliferation, metabolism and apoptosis (3). Endogenous sources of free radicals include the mitochondrial respiratory chain, inflammation, peroxisomes and cytochrome P450 (4). However, excessive production of ROS can lead to oxidative stress, a pathophysiological condition, resulting in oxidative damage of macromolecules (lipids, proteins and DNA) (1, 5). Thus, oxidative stress has been associated with several pathophysiological conditions and diseases (1).

One of the pathological conditions associated with oxidative stress is sepsis, a systemic inflammatory response syndrome attributed to infection (6-8). The infection is most commonly caused by bacteria but can also be caused by fungi, viruses, or parasites (6, 7). Common symptoms of sepsis include fever, increased heart rate, increased breathing rate, and confusion (6, 7). Sepsis leads to collateral damage of normal tissues due to the constant activation of monocytes and endothelial cells that trigger inflammatory and coagulation cascades (9). The severity and outcome of septic syndrome largely depend on the original site and type of infection and the host's inflammatory response, as well as the time of administration of appropriate antimicrobial therapy (10-12).

Another pathological condition that has been associated with oxidative stress is obesity (13,14). Obesity is a chronic metabolic disease in which excess body fat in adipose tissue has accumulated to the extent that it may have a negative effect on health, leading to reduced life expectancy and increased health problems such as heart disease, type 2 diabetes, obstructive sleep apnea, certain types of cancer, and osteoarthritis (14, 15). Since increased oxidative stress is observed in obese individuals, examination of the impact of obesity on sepsis-induced oxidative stress would be interesting (13, 14).

Correspondence to: Dimitrios Kouretas, Department of Biochemistry and Biotechnology, University of Thessaly, Ploutonos 26 & Aioulou, Larissa 41221, Greece. Tel: +30 2410565277, Fax: +30 2410565293, e-mail: dkouret@uth.gr

Key Words: Oxidative stress, sepsis, obesity, biomarkers, oxidation-reduction potential.

A number of biomarkers have been developed for assessing oxidative stress levels in humans. These biomarkers help to assess the redox status of a person and consequently predict the possible manifestation of a pathological condition or the progress of an already present disease (16). However, oxidative stress measurement is still incomplete, time consuming and impractical in a clinical setting, and so there is a great need for developing new methods for assessing the redox status in humans (17). A novel method for assessing oxidative stress is the proprietary RedoxSYS Diagnostic System (Luoxis Diagnostics, Inc., Englewood, CO, USA). The RedoxSYS Diagnostic System measures the oxidation-reduction potential (ORP), which is an integrated measure of the balance between total oxidants (*e.g.* oxidized thiols, superoxide radical, hydroxyl radical, hydrogen peroxide, nitric oxide, peroxyxynitrite and transition metal ions), and total reductants (*e.g.* free thiols, ascorbate, α -tocopherol, β -carotene and uric acid). Thus, ORP is an overall measure of the oxidative stress to which a biological system is subjected (18). The RedoxSYS Diagnostic System enables robust and rapid assessment of oxidative stress in a single drop of plasma *via* measurement within four minutes of two distinct elements to determine ORP, namely the static ORP (sORP) and the capacity ORP (cORP). sORP is the standard potential between a working electrode and a reference electrode with no driving current (or extremely small current) which is proportional to the balance of reductants and oxidants and is what is classically termed ORP (*i.e.* a homeostatic parameter capturing the current balance of oxidants and reductants in a biological specimen). Low sORP values mean that the biological sample is in the normal range of oxidative stress, while higher than normal sORP values means that the biological sample is in a higher state of oxidative stress. cORP is the measure of antioxidant reserve available in the body's system; high capacity values mean that the biological sample has antioxidant reserves in the normal range; lower than normal cORP values means that the biological sample has below normal antioxidant reserves.

In previous studies, we have shown that ORP markers could be used for assessing oxidative stress induced by physiological conditions such as exercise (19-20). In the present study, ORP markers were used for the first time for assessing oxidative stress levels in a pathological condition such as sepsis. The possible effects of obesity-induced oxidative stress on patients with sepsis were also investigated. Apart from ORP markers, conventional oxidative stress biomarkers were used such as total antioxidant capacity (TAC), thiobarbituric acid-reactive substances (TBARS) and protein carbonyls (CARB).

Materials and Methods

Participants. In this prospective study, a total of 42 individuals who were admitted to Patras University General Hospital in Greece were

enrolled. According to our protocol, the patients were divided into four groups. The inclusion criteria of each group were: Control group (n=11): individuals with body mass index (BMI) <30 kg/m² without clinicolaboratory signs of infection; Obesity group (n=10): individuals with BMI \geq 30 kg/m² without clinicolaboratory signs of infection; Sepsis group (n=12): individuals with BMI <30 kg/m² with sepsis; Sepsis and obesity group (n=9): individuals with BMI \geq 30 kg/m² with sepsis. Sepsis was defined according to the criteria of the American College of Chest Physicians–Society of Critical Care Medicine Consensus Conference Committee—as the presence of confirmed infection and >2 of the following criteria: (a) a temperature of >38°C or <36°C; (b) a heart rate of >90 beats per min; (c) tachypnea, manifested by a respiratory rate of >20 breaths per min or hyperventilation, indicated by a PaCO₂ of <32 mm Hg; and (d) an altered white blood cell count of >12000 or <4000 cells per mm³ or the presence of >10% immature forms.

Blood collection and handling. Blood was collected in ethylenediamine tetra-acetic acid tubes for measuring oxidative stress markers. Blood samples were centrifuged immediately at 1370 \times g for 10 min at 4°C and the plasma was collected and used for the measurements of oxidative stress markers. Plasma samples were stored at –80°C prior to biochemical analyses.

Assessment of sORP and cORP using the the RedoxSYS Diagnostic System. sORP and cORP values were determined using the RedoxSYS Diagnostic System (Luoxis Diagnostics, Inc., Englewood, CO, USA). Twenty microliters of plasma was applied to disposable sensors designed by Luoxis, which were inserted into the RedoxSYS Diagnostic System that measured and reported within four minutes the sORP and cORP values. The RedoxSYS diagnostic system measures the ORP with a three-electrode system: a working electrode, a counter electrode and a reference electrode. Firstly, a negligible amount of current is applied between the working and counter electrodes, and the ORP is measured between them. Once the ORP reading reaches equilibrium, the sORP is established and measured in millivolts. Then a linearly increasing current is applied to the sample, between the counter and working electrodes. The time from the beginning of the current sweep to the maximum rate of change in ORP is referred to as the transition time and the integrated current to this time is the cORP, measured in microcoulombs. Each sample was measured in triplicate.

Assessment of TAC, TBARS and CARB. For TBARS determination, a slightly modified assay of Keles *et al.* was used (21). According to this method, 100 μ l of plasma was mixed with 500 μ l of 35% trichloroacetic acid (TCA) and 500 μ l of tris(hydroxymethyl)aminomethane hydrochloride (Tris-HCl) (200 mM, pH 7.4) and incubated for 10 min at room temperature. One milliliter of 2 M Na₂SO₄ and 55 mM thiobarbituric acid solution was added and the samples were incubated at 95°C for 45 min. The samples were cooled on ice for 5 min and were vortexed after adding 1 ml of 70% TCA. The samples were centrifuged at 15,000 \times g for 3 min and the absorbance of the supernatant was read at 530 nm. A baseline absorbance was taken into account by running a blank along with all samples during the measurement. Calculation of TBARS concentration was based on the molar extinction coefficient of malon dialdehyde.

The determination of CARB was based on the method of Patsoukis *et al.*, (22). The derivatization of the carbonyl group with

2,4-dinitrophenylhydrazine (DNPH), which leads to the formation of a stable 2,4-dinitrophenyl (DNP) hydrazone product, is one of the most widely utilized measurements of protein oxidation (23). It has been shown to be very sensitive, especially for the determination of carbonyl content in purified proteins (23). For this reason, the proteins were precipitated with TCA. In this assay, 50 μ l of 20% TCA was added to 50 μ l of plasma and this mixture was incubated in an ice-bath for 15 min and centrifuged at $15,000 \times g$ for 5 min at 4°C . The supernatant was discarded and 500 μ l of 10 mM DNPH (in 2.5 N HCl) for the sample, or 500 μ l of 2.5 N HCl for the blank, was added to the pellet. The samples were incubated in the dark at room temperature for 1 h, with intermittent vortexing every 15 min and were centrifuged at $15,000 \times g$ for 5 min at 4°C . The supernatant was discarded and 1 ml of 10% TCA was added, vortexed and centrifuged at $15,000 \times g$ for 5 min at 4°C . The supernatant was discarded and 1 ml of ethanol-ethyl acetate (1:1 v/v) was added, vortexed and centrifuged at $15,000 \times g$ for 5 min at 4°C . This washing step was repeated twice. The supernatant was discarded and 1 ml of 5 M urea (pH 2.3) was added, vortexed and incubated at 37°C for 15 min. The samples were centrifuged at $15,000 \times g$ for 3 min at 4°C and the absorbance was read at 375 nm. Calculation of CARB concentration was based on the molar extinction coefficient of DNPH. Total plasma protein was assayed using Bradford reagent from Sigma-Aldrich (Munich, Germany).

The determination of TAC was based on the method of Janaszewska and Bartosz (24). Briefly, 20 μ l of plasma were added to 480 μ l of 10 mM sodium potassium phosphate (pH 7.4) and 500 μ l of 0.1 mM 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical and the samples were incubated in the dark for 30 min at room temperature. The samples were centrifuged for 3 min at $20,000 \times g$ and the absorbance was read at 520 nm. TAC is presented as mmol of DPPH reduced to 2,2-diphenyl-1-picrylhydrazine (DPPH:H) by plasma antioxidants.

Statistical analysis. For statistical analysis, data were analyzed by one-way ANOVA followed by Dunnett's test for multiple pair-wise comparisons. The level of statistical significance was set at $p < 0.05$. For all statistical analyses, SPSS, version 13.0 (SPSS Inc., Chicago, IL, USA) was used. Data are presented as the mean \pm SEM.

Results

From the two ORP parameters measured by the RedoxSYS Diagnostic System, the sORP indicating the current redox balance was increased significantly ($p < 0.05$) by 9.3% in the sepsis group, by 6.9% in the sepsis and obesity group compared to their control group (Figure 1A). On the contrary, cORP, reflecting the reserves of antioxidants, was significantly lower by 11.8% in the sepsis group by 17.7% in the sepsis and obesity group compared to control group (Figure 1B). Neither of these two markers differed significantly in the obesity group compared to the control group (Figures 1A and B).

Regarding the conventional oxidative stress markers, the plasma levels of CARB indicating protein oxidation were increased significantly ($p < 0.05$) by 73.8% in the obesity group and by 167.5% in the sepsis and obesity group compared to the controls, while there was not any significant




Figure 1. Oxidation reduction potential (ORP) parameters of the redox status in plasma of control group, sepsis group, obesity group, and obesity and sepsis group as measured by the RedoxSYS Diagnostic System. sORP, Static oxidation-reduction potential; cORP, capacity oxidation-reduction potential. *Significantly different compared to the control group ($p < 0.05$).

change in the sepsis group. Moreover, in the sepsis and obesity group, CARB levels were increased significantly by 130.1% and 53.9% ($p < 0.05$) compared to the sepsis group and obesity group, respectively (Figure 2A). TAC was significantly ($p < 0.05$) lower by 20.6% in the sepsis group and by 30.8% in the sepsis and obesity group compared to the control group, while there was no significant difference




Figure 2. Conventional parameters of redox status in plasma of the control group, sepsis group, obesity group, and obesity and sepsis group. A: Protein carbonyl levels (CARB); significantly different at $p < 0.05$ compared to *control group, and #sepsis group and obesity group. B: Total antioxidant capacity (TAC); significantly different at $p < 0.05$ compared to *control group, and #obesity group. C: Thiobarbituric acid-reactive substances (TBARS); significantly different at $p < 0.05$ compared to *control group, and #sepsis group.

comparing controls with the obesity group (Figure 2B). In the sepsis and sepsis and obesity groups, TAC was significantly ($p < 0.05$) lower by 23.4% and 33.3%, respectively, compared to the obesity group (Figure 2B). TBARS, reflecting lipid peroxidation, were significantly ($p < 0.05$) higher by 59.1% in the obesity group and by 83.6% in the sepsis and obesity group compared to controls, while there was not any significant difference for the sepsis group (Figure 2C). In the obesity group, and sepsis and obesity group, the levels of TBARS were significantly higher by 42.8% and 64.8%, respectively ($p < 0.05$), compared to the sepsis group (Figure 2C).

Discussion

Sepsis is defined as the systemic inflammatory response syndrome attributed to infection (6, 7). Several studies have reported higher levels of oxidative stress and lower antioxidant potential in patients with sepsis in the first hours

or days after ICU admission compared to healthy controls (25, 26). In the present study, we examined if ORP markers as measured by the RedoxSYS Diagnostic System could be used for assessing oxidative stress in patients with sepsis. The results suggested that both sORP and cORP can detect sepsis-induced oxidative stress. Specifically, sepsis increased sORP, reflecting the overall current redox balance, thus higher than normal sORP values means that the biological sample is in a higher state of oxidative stress. Moreover, sepsis reduced cORP indicating a reduced antioxidant reserve. It has been proposed that sepsis-induced oxidative stress is due to the continuous defense and inflammatory response during sepsis that results in a large increase in the levels of free radicals, nitric oxide and inflammatory cytokines (27).

Among the conventional oxidative stress markers, only TAC indicated sepsis-induced oxidative stress, it was lower in patients with sepsis compared to control individuals. On the contrary, there were no significant changes in CARB and TBARS between patients with sepsis and control individuals. These results for the conventional biomarkers were in agreement with one of our previous studies in which oxidative stress was assessed in patients with sepsis (28). However, in another study in which the variability of oxidative stress during sepsis evolution was investigated, we found that severe sepsis resulting in mortality increased CARB in plasma, while it had no effect on TAC and TBARS levels (29). This latter study demonstrated different effects of sepsis on TAC and TBARS than were found in the present study. Obviously, an explanation for this inconsistency may be the different severity of the sepsis manifested by the patients of the two studies. On the other hand, we believe that this inconsistency may emphasize the need for finding new markers that would more consistently assess oxidative stress in sepsis and especially would determine an overall redox status of an individual. Most of the current oxidative stress markers assess a single parameter of oxidative damage (e.g. lipid peroxidation, protein oxidation) or of antioxidant defense (e.g. glutathione GSH levels, catalase activity, superoxide dismutase activity). Consequently, several of these parameters should be measured in order to assess oxidative stress more reliably in an individual. However, the measurement of several biomarkers of oxidative stress is time consuming, and is impractical in a clinical setting such as in the case of sepsis. Thus, the finding that ORP markers as evaluated by the RedoxSYS Diagnostic System can consistently measure sepsis-induced oxidative stress is important, since they are measured quickly (within four minutes) and easily. Moreover, ORP markers are an integrated oxidative stress measurement, since they assess the total imbalance between oxidants and reductants (i.e. sORP) or the total antioxidant defence (i.e. cORP). Thus, the RedoxSYS Diagnostic System could be used at the bedside for monitoring oxidative stress in patients with sepsis similar

to the bedside monitoring of heart rate, respiration, *etc.*

ORP markers were used to assess the effects of obesity on sepsis-induced oxidative stress, since obesity alone induces oxidative stress (30). Obesity-induced oxidative stress has been attributed to hyperglycemia, increased muscle activity in order to carry excessive weight, elevated tissue lipid levels and low-grade endotoxemia due to gut barrier dysfunction (30). In a previous study, we reported that obesity was associated with increased oxidative stress in patients with sepsis (28). The present results from the conventional oxidative stress biomarkers confirmed those of our previous study, that is, obesity increased lipid oxidation (*i.e.* TBARS) and protein oxidation (*i.e.* CARB) in patients with sepsis. These effects were probably due to the combination of the factors inducing oxidative stress in each of the two conditions, sepsis and obesity. However, ORP markers did not detect increased oxidative stress in obese patients with sepsis compared to patients with sepsis alone. Moreover, ORP markers did not reveal increased oxidative stress in obese individuals compared to controls. Like ORP markers, TAC, the other holistic marker measuring total antioxidant defense, did not increase in obese individuals compared to controls, while obese individuals had higher protein oxidation and lipid peroxidation. Other studies have shown that obesity reduced TAC, but these were large studies including more than 1500 individuals (31). Thus, the fact that ORP and TAC markers did not detect differences in oxidative stress between nonobese and obese individuals may be due to the small number of participants in our study. We are currently performing a study including more than 500 individuals in order to examine the assessment of oxidative stress using ORP markers in metabolic syndromes such as obesity.

In conclusion, the present results suggest for the first time that ORP markers as measured by the RedoxSYS Diagnostic System could be used for assessing oxidative stress in patients with sepsis. Currently, we are carrying out a study to examine if ORP markers correlate with injury severity and degree of inflammation in patients with sepsis. The main advantage of using sORP and cORP markers for measuring oxidative stress is their rapid measurement, using a small plasma volume (20 μ l), of the total imbalance between oxidants and reductants, and the total antioxidant defense levels respectively. Thus, these markers could be used for alerting a physician to the oxidative stress levels in patients with sepsis so that appropriate intervention (*e.g.* antioxidant administration) can be applied (32). Importantly, studies have shown that the timely administration of antioxidant is more effective in critically ill patients than delayed treatment (33).

References

- Halliwell B: The wanderings of a free radical. *Free Radic Biol Med* 46(5): 531-542, 2009.

- Dupre-Crochet S, Erard M and Nubetae O: ROS production in phagocytes: Why, when, and where? *J Leukoc Biol* 94(4): 657-670, 2013.
- Sinha K, Das J, Pal PB and Sil PC: Oxidative stress: the mitochondria-dependent and mitochondria-independent pathways of apoptosis. *Arch Toxicol* 2013 87(7): 1157-1180, 2013.
- Valko, M, Leibfritz D, Moncol J, Cronin MTD, Mazur M and Telser J: Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 39(1): 44-84, 2007.
- Mylonas C and Kouretas D: Lipid peroxidation and tissue damage. *In Vivo* 13(3): 295-309, 1999.
- Deutschman CS and Tracey KJ: Sepsis: Current dogma and new perspectives. *Immunity* 40(4): 463-75, 2014.
- Green JM: Essentials of Sepsis Management. *Surg Clin North Am* 95(2): 355-365, 2015.
- David BO, Carrick MM, Mains CW, Rael LT, Slone D and Brody EN: Sepsis, oxidative stress, and hypoxia: Are there clues to better treatment? *Redox Rep* (in press), 2015.
- Remick DG: Pathophysiology of Sepsis. *Am J Pathol* 170(5): 1435-1444, 2007.
- Hall TC, Bilku DK, Al-Leswas D, Horst C and Dennison AR: Biomarkers for the differentiation of sepsis and SIRS: the need for the standardisation of diagnostic studies. *Ir J Med Sci* 180(4): 793-798, 2007.
- Martin GS: Sepsis, severe sepsis and septic shock: changes in incidence, pathogens and outcomes. *Expert Rev Anti Infect Ther* 10(6): 701-706, 2012.
- Perman SM, Goyal M and Gaieski DF: Initial emergency department diagnosis and management of adult patients with severe sepsis and septic shock. *Scand J Trauma Resusc Emerg Med* 20: 41, 2012.
- Matsuda M and Shimomura I: Increased oxidative stress in obesity: implications for metabolic syndrome, diabetes, hypertension, dyslipidemia, atherosclerosis, and cancer. *Obes Res Clin Pract* 7(5): e330-341, 2013.
- Rentería I, Arenas Berumen E, Arellano García ME, Carrasco-Legleu CE, De León-Fierro LG and Arenas-Berumen EA: Factors affecting oxidative damage in obese children: an exploratory study. *Nutr Hosp* 31(4): 1499-1503, 2015.
- Haidar YM and Cosman BC: Obesity epidemiology. *Clin Colon Rectal Surg* 24(4): 205-210, 2011.
- Dalle-Donne I, Rossi R, Colombo R, Giustarini D and Milzani A: Biomarkers of oxidative damage in human disease. *Clin Chem* 52(4): 601-623, 2006.
- Ogino K and Wang DH: Biomarkers of oxidative/nitrosative stress: an approach to disease prevention. *Acta Med Okayama* 61(4): 181-189, 2007.
- Harris C and Hansen JM: Oxidative stress, thiols, and redox profiles. *Methods Mol Biol* 889: 325-346, 2012.
- Stagos D, Goutzourelas N, Bar-Or D, Ntontou AM, Bella E, Becker AT, Statiri A, Kafantaris I and Kouretas D: Application of a new oxidation-reduction potential assessment method in strenuous exercise-induced oxidative stress. *Redox Rep* (in press), 2014.
- Stagos D, Goutzourelas N, Ntontou A-M, Kafantaris I, Deli CK, Poullos A, Jamurtas AZ, Bar-Or B and Dimitrios Kouretas D: Assessment of eccentric exercise-induced oxidative stress using oxidation-reduction potential markers. *Oxid Med Cell Longev* 2015: 204615, 2015.

- 21 Keles MS, Taysi S, Sen N, Aksoy H and Akcay F: Effect of corticosteroid therapy on serum and CSF malondialdehyde and antioxidant proteins in multiple sclerosis. *Can J Neurol Sci* 28(2): 141-143, 2001.
- 22 Patsoukis N, Zervoudakis G, Panagopoulos NT, Georgiou CD, Angelatou F and Matsokis NA: Thiol redox state (TRS) and oxidative stress in the mouse hippocampus after pentylene-tetrazol-induced epileptic seizure. *Neurosci Lett* 357(2): 83-86, 2004.
- 23 Luo S and Wehr NB: Protein carbonylation: avoiding pitfalls in the 2,4-dinitrophenylhydrazine assay. *Redox Rep* 14(4): 159-166, 2009.
- 24 Janaszewska A and Bartos G: Assay of total antioxidant capacity: comparison of four methods as applied to human blood plasma. *Scandinavian J Clin Lab Invest* 62(3): 231-236, 2002.
- 25 Abiles J, de la Cruz AP, Castaño J, Rodríguez-Elvira M, Aguayo E, Moreno-Torres R, Llopis J, Aranda P, Argóelles S, Ayala A, de la Quintana AM and Planells EM: Oxidative stress is increased in critically ill patients according to antioxidant vitamin intake, independent of severity: a cohort study. *Crit Care* 10(5): R146, 2006.
- 26 Alonso de Vega J, Dvaz J, Serrano E and Carbonell L: Oxidative stress in critically ill patients with systemic inflammatory response syndrome. *Crit Care Med* 30: 1782-1785, 2002.
- 27 Victor VM, Espulgues JV, Hernandez-Mijares A and Rocha M: Oxidative stress and mitochondrial dysfunction in sepsis: a potential therapy with mitochondria-targeted antioxidants. *Infect Disord Drug Targets* 9(4): 376-389, 2009.
- 28 Kolyva AS, Zolota V, Mpatoulis D, Skroubis G, Solomou EE, Habeos IG, Assimakopoulos SF, Goutzourelas N, Kouretas D and Gogos CA: The role of obesity in the immune response during sepsis. *Nutr Diabetes* 4: e137, 2014.
- 29 Karapetsa M, Pitsika M, Goutzourelas N, Stagos D, Tousia Becker A and Zakynthinos E: Oxidative status in ICU patients with septic shock. *Food Chem Toxicol* 61: 106-111, 2013.
- 30 Vincent HK and Taylor AG: Biomarkers and potential mechanisms of obesity-induced oxidant stress in humans. *Int J Obes* 30(3): 400-418, 2006.
- 31 Chrysohoou C, Panagiotakos DB, Pitsavos C, Skoumas I, Papademetriou L, Economou M and Stefanadis C: The implication of obesity on total antioxidant capacity in apparently healthy men and women: the ATTICA study. *Nutr Metab Cardiovasc Dis* 17(8): 590-597, 2007.
- 32 Mishra V: Oxidative stress and role of antioxidant supplementation in critical illness. *Clin Lab* 53(3-4): 199-209, 2007.
- 33 Lovat R and Preiser JC: Antioxidant therapy in intensive care. *Curr Opin Crit Care* 9(4): 266-270, 2003.

Received April 20, 2015

Revised June 2, 2015

Accepted June 4, 2015

June 4, 2015

Dr. Dimitrios Kouretas

Re: Your manuscript No. **3264-S** entitled «**Assessment of Oxidative Stress...**»

Dear Dr

Referring to your above manuscript for publication in *in vivo*, please allow us to use this form letter in reply:

1. *Referee's recommendations:*

- Urgent to be published immediately.
- Accepted in the presented form.
- Accepted with minor changes.
- Accepted with grammatical or language corrections.
- Remarks: **Please provide Figures at a higher quality resolution**

2. *Excess page charges.*

- Your article has approx. **xxx** printed pages and is in excess of the allotted number by approx. **xx** printed pages. The charges are EURO € **xxx** per excess page, totalling EURO € **xxx**
We ask you to confirm acceptance of these charges.
- Your article includes **xxx** pages with color figures. The charges are EURO € **xxx** per color page, totalling € EURO **xxxx**
- Our invoice will be sent by e-mail to the corresponding author.

3. Your article will appear in Volume **29**, Issue No. **5**, **2015**

4. Please order your reprints, pdf or online open access, now. This will facilitate our prompt planning of future issues and rapid publication of your article. Reprints will be delivered by rapid one-day delivery within one month from publication.

We would appreciate your prompt reply.

With many thanks,

Yours sincerely,



J.G. Delinasios

Managing Editor

EDITORIAL OFFICE: INTERNATIONAL INSTITUTE OF ANTICANCER RESEARCH

DELINASIOS G.J. & CO G.P., Kapandriti, P.O.B. 22, Attiki 19014, Greece. Tel.: 0030-22950-52945;

Tel & Fax:0030-22950-53389; e-mail: journals@iiar-anticancer.org

Editorial Office: International Institute of Anticancer Research,
DELINASIOS G.J. & CO G.P., Kapandriti, P.O.B. 22, Attiki 19014, Greece
Fax: +30-22950-53389; Tel: +30-22950-52945; e-mail: journals@iiar-anticancer.org

Please type or print the requested information on the reprint order form and return it to the Editorial Office by fax or e-mail. Fees for reprints, PDF file, or online open access must be paid for in advance.

If your paper is subject to charges for excess pages or color plates, please add these charges to the payment for reprints. The reprints are not to be sold.

PRICE LIST FOR REPRINTS WITHOUT COVER

Page length	Online Open Access Fee* (Euro)	PDF File Fee (Euro)	Number of copies requested (prices are in Euro)									
			50	100	200	300	400	500	1000	1500	2000	3000
1-4pp	400	175	140	285	337	388	453	504	851	1135	1470	2038
5-8	500	225	150	388	453	530	595	672	1083	1445	1832	2554
9-12	600	277	160	504	569	659	737	827	1341	1780	2219	3096
13-16	700	354	180	659	737	840	943	1046	1625	2141	2657	3676
17-20	800	419	200	788	879	982	1098	1227	1883	2451	3044	4244

***Online open access of an article published in 2015 is accompanied by a complimentary online subscription to *in vivo*.**

For reprints with cover: Please add EURO 80.00 per 100 copies.

Postage: Please add 5% on the reprint prices.

Reprint Order Form

Of my paper No. **3264-S** comprising **6** printed pages, entitled «**Assessment of Oxidative Stress...**» accepted for publication in **in vivo** Vol. **29** No. **5**

I require a total of _____ copies at EURO:

I do not require reprints.

Please send me a PDF file of the article at EURO:

Please provide Online Open Access of the article at the Stanford University Highwire Press website (at Euro _____) immediately upon publication, and enter my complimentary online subscription to **in vivo**.

Please send me a copy of the issue containing my paper at EURO 25.00.

Please enter my personal subscription to **in vivo** at the special Author's price of EURO 190.00 (print; online) (Year: 2015).

A check for the above amounts payable to DELINASIOS G.J. & CO G.P., is enclosed.

Please send an invoice to (Billing Name and Address):

VAT number (For EC countries):

Name:

Address:

Country:

Postal code:

Tel:

City:

e-mail:

Fax:

Signature:

Please send reprints to (Complete Address and Tel. no.):